Version: 0 February 14 2014

Composition of Two Potentials

Khosrow Chadan[1], Reido Kobayashi[2] and Ryu Sasaki[3]

[1] Laboratoire de Physique Théorique*
Université de Paris XI, Bâtiment 210, 91405 Orsay Cedex, France
[2] Department of Mathematics
Tokyo University of Science, Noda, Chiba 278-8510, Japan
[3] Department of Physics
Shinshu University, Matsumoto 390-8621, Japan

arXiv:1402.6954v1 [math-ph] 27 Feb 2014

Abstract

Given two potentials V_0 and V_1 together with a *certain nodeless solution* φ_0 of V_0, we form a composition of these two potentials. If V_1 is exactly solvable, the composition is exactly solvable, too. By combining various solvable potentials in one-dimensional quantum mechanics, a huge variety of solvable compositions can be made.

1 Introduction

Some years ago, two of the present authors developed a method of composing two potentials V_0 and V_1 in a special setting [1]. The potentials were spherically symmetric and of short range and were explicitly solvable at zero energy. The starting potential V_0 should sustain no bound states. Then the composed potential was explicitly solvable at zero energy, too. The main motivation was applications to quantum mechanical scattering problems [2] and the radial Sch"oinger equations.

In the present paper, we show that the composition of two potentials V_0 and V_1 in generic one dimensional quantum mechanics can be defined in a much more general setting. The only requirement is that V_0 should have a *nodeless solution* satisfying certain boundary conditions. If V_1 is exactly solvable, so is the composed potential V_C. If V_1 has also a nodeless solution satisfying boundary conditions, a further composition with another potential V_2 can be made, as emphasised in the earlier papers [1]. We provide several examples of V_0,

*Unité Mixe de Recherche UMR 8627-CRNS

all exactly solvable, together with an infinite number of nodeless solutions satisfying the boundary conditions. These nodeless solutions are called *virtual state wavefunctions* [3, 4], which have played an important role in the construction of various rational deformations of exactly solvable potentials and the *multi-indexed orthogonal polynomials* [3]–[12] as the main part of the eigenfunctions of the deformed solvable systems.

This paper is organised as follows. In section 2, we formulate the composition of two potentials in the most general setting. Derivation is quite elementary. Section 3 provides explicit examples of nodeless solutions (virtual state wavefunctions) belonging to the well-known solvable potentials [13, 14]; the radial oscillator potential in §3.1, Pöschl-Teller potential in §3.2, hyperbolic Pöschl-Teller potential in §3.3, Rosen-Morse potential in §3.4 and Eckart potential in §3.5. The final section is for a short summary and comments.

2 General Composition

The main ingredients of the present theory are two quantum mechanical systems in one dimension with smooth potentials V_0 and V_1:

$$\text{System 0}: \qquad \mathcal{H}_0 = -\frac{d^2}{dx^2} + V_0(x), \qquad a < x < b, \tag{2.1}$$

$$\mathcal{H}_0\varphi_0(x) = \tilde{\mathcal{E}}_0\varphi_0(x), \tag{2.2}$$

$$\text{System 1}: \qquad \mathcal{H}_1 = -\frac{d^2}{dx^2} + V_1(x), \qquad c < x < d, \tag{2.3}$$

$$\mathcal{H}_1\phi_m(x) = \mathcal{E}_m\phi_m(x), \qquad m = 1, \ldots, \tag{2.4}$$

Since they are both quantum mechanical systems, a finite boundary can only be realised by a strong repulsive singularity. Otherwise, the wave functions would leak beyond that boundary and the quantum system is ill-defined. The System 0 needs not be solvable. The only requirement is that it has a *nodeless real solution* $\varphi_0(x)$ with proper boundary conditions (A) and (B), to be specified later. The System 0 may or may not have some discrete eigenstates, either finitely many or infinite in number. The System 1 is a general quantum system. Nothing special is required of $V_1(x)$ for the construction of the composition of the two potentials V_0 and V_1.

It is elementary to show that the following function

$$\chi_0(x) \stackrel{\text{def}}{=} \varphi_0(x)\left(\alpha + \int_x^b \frac{dt}{\varphi_0^2(t)}\right), \quad \alpha \stackrel{\text{def}}{=} \frac{1}{d-c}, \tag{2.5}$$

is another solution of $\mathcal{H}_0$ with the same energy $\tilde{\mathcal{E}}_0$:

$$\mathcal{H}_0\chi_0(x) = \tilde{\mathcal{E}}_0\chi_0(x). \tag{2.6}$$

Let us introduce a mapping function $\psi_0(x)$, $a < x < b$,

$$\psi_0(x) \stackrel{\text{def}}{=} c + \frac{\varphi_0(x)}{\chi_0(x)} = c + \frac{1}{\alpha + \int_x^b \frac{dt}{\varphi_0^2(t)}}, \tag{2.7}$$

which is monotonously increasing

$$\frac{d\psi_0(x)}{dx} = \frac{1}{\chi_0^2(x)} > 0.$$

We assume the following boundary conditions on $\varphi_0(x)$:

$$x \to a \qquad \int_x^b \frac{dt}{\varphi_0^2(t)} \to \infty \tag{A},$$

$$x \to b \qquad \int_x^b \frac{dt}{\varphi_0^2(t)} \to 0 \tag{B},$$

which mean

$$\psi_0(a) = c, \quad \psi_0(b) = c + \frac{1}{\alpha} = d. \tag{2.8}$$

Thus $\psi_0(x)$ is a 1:1 onto mapping from (a, b) to (c, d).

The composition of two potentials, V_0 and V_1 is achieved by $\chi_0(x)$ and $\psi_0(x)$:

$$\mathcal{H}_C = -\frac{d^2}{dx^2} + V_C(x), \quad V_C(x) \stackrel{\text{def}}{=} V_0(x) - \tilde{\mathcal{E}}_0 + \frac{1}{\chi_0^4(x)} V_1(\psi_0(x)), \quad a < x < b. \tag{2.9}$$

With a solution $\phi_m(x)$ of $\mathcal{H}_1$, let us define

$$\phi_m^C(x) \stackrel{\text{def}}{=} \chi_0(x)\phi_m(\psi_0(x)), \quad m = 1, \ldots, . \tag{2.10}$$

It is elementary to show that $\phi_m^C(x)$ satisfies

$$\mathcal{H}_C\phi_m^C(x) = \frac{\mathcal{E}_m}{\chi_0^4(x)}\phi_m^C(x), \quad m = 1, \ldots. \tag{2.11}$$

If $\{\phi_m(x)\}$ are eigenfunctions of $\mathcal{H}_1$:

$$\int_c^d \phi_m(x)\phi_n(x)dx = h_m\delta_{m\,n}, \quad h_m > 0, \quad m, n = 1, \ldots, \tag{2.12}$$

then $\{\phi_m^C(x)\}$ are orthogonal with each other with respect to the weight function $\frac{1}{\chi_0^4(x)}$:

$$\int_a^b \phi_m^C(x)\phi_n^C(x)\frac{dx}{\chi_0^4(x)} = \int_a^b \phi_m(\psi_0(x))\phi_n(\psi_0(x))\frac{dx}{\chi_0^2(x)} \tag{2.13}$$

$$= \int_c^d \phi_m(\psi_0)\phi_n(\psi_0)d\psi_0 = h_m\delta_{m\,n}, \quad m,n = 1,\ldots,. \tag{2.14}$$

If $\mathcal{H}_1$ is exactly solvable, so is the composed Hamiltonian $\mathcal{H}_C$.

As stressed in [1], further compositions are straightforward, so long as the proper nodeless solutions of can be found. Suppose we have another Hamiltonian system $\mathcal{H}_2$ with a potential V_2

$$\begin{aligned} \text{System 2}: \qquad \mathcal{H}_2 &= -\frac{d^2}{dx^2} + V_2(x), \qquad && e < x < f, \\ \mathcal{H}_2\Phi_m(x) &= E_m\Phi_m(x), && m = 1,\ldots, \end{aligned}$$

and a nodeless solution φ_1 of $\mathcal{H}_1$ with proper boundary conditions. The next step is essentially the same as before:

$$\begin{aligned} &\mathcal{H}_1\varphi_1(x) = \tilde{\mathcal{E}}_1\varphi_1(x), \quad \chi_1(x) \stackrel{\text{def}}{=} \varphi_1(x)\left(\alpha_1 + \int_x^d \frac{dt}{\varphi_1^2(t)}\right), \quad \alpha_1 = (f-e)^{-1}, \\ &\mathcal{H}_1\chi_1(x) = \tilde{\mathcal{E}}_1\chi_1(x), \quad \psi_1(x) \stackrel{\text{def}}{=} e + \frac{\varphi_1(x)}{\chi_1(x)} = e + \frac{1}{\alpha_1 + \int_x^d \frac{dt}{\varphi_1^2(t)}}, \quad \frac{d\psi_1(x)}{dx} = \frac{1}{\chi_1^2(x)} > 0. \end{aligned}$$

The second composed Hamiltonian is

$$\begin{aligned} \mathcal{H}_{C2} &= -\frac{d^2}{dx^2} + V_{C2}(x), \qquad && a < x < b, \\ V_{C2}(x) &\stackrel{\text{def}}{=} V_0(x) - \tilde{\mathcal{E}}_0 + \frac{1}{\chi_0^4(x)}\Big(V_1(\psi_0(x)) - \tilde{\mathcal{E}}_1\Big) + \frac{1}{\chi_0^4(x)\chi_1^4(\psi_0(x))}V_2\big(\psi_1(\psi_0(x))\big), \end{aligned}$$

with its solutions

$$\begin{aligned} \Phi_m^{C2}(x) &\stackrel{\text{def}}{=} \chi_0(x)\chi_1(\psi_0(x))\Phi_m\big(\psi_1(\psi_0(x))\big), \quad m = 1,\ldots, \\ \mathcal{H}_{C2}\Phi_m^{C2}(x) &= \frac{E_m}{\chi_0^4(x)\chi_1^4(\psi_0(x))}\Phi_m^{C2}(x). \end{aligned}$$

The composition processes can go on indefinitely.

3 Explicit Examples

Several explicit examples of the compositions were demonstrated in [1] in connection with certain scattering problems [2]. For example, given two spherically symmetric and short range potentials for which the radial Schrödinger equations can be solved at zero energy, the composition can also be solved at zero energy. Here we give several explicit examples of the System 0, together with the proper *nodeless solutions* $\varphi_0(x)$. Those examples presented in

[1] will not be re-listed here. Although System 0 needs not be solvable, most examples listed below are, in fact, exactly solvable. The reason is quite trivial. The proper nodeless solutions can be most easily found when the system is solvable. The nodeless solutions listed below are called *virtual state wavefunctions* [3, 4]. Their energies are *below the ground state energy* and they together with their inverses are *square non-integrable*. Thus the conditions (A) and (B) are satisfied. In most cases, they are obtained from the eigenfunctions by discrete symmetry transformations. They have played an essential role in the rational deformations of solvable potentials, creating the multi-indexed and exceptional orthogonal polynomials [3], [5]–[12], which are new species of orthogonal polynomials satisfying second order differential equations but not the three term recurrence relations [15].

See for example [13, 14] for a general review of exactly solvable potentials. We follow [13] for the naming of solvable potentials. Detailed information of these systems, the symmetry, eigenfunctions, various virtual state wave functions and their generalisation can be found in [3, 4]. The first two examples have infinitely many discrete eigenstates.

3.1 Radial oscillator

The radial oscillator potential is

$$V_0(x) \stackrel{\text{def}}{=} x^2 + \frac{g(g-1)}{x^2}, \quad 0 < x < \infty, \quad g > 1/2. \tag{3.1}$$

The lower boundary $x = 0$ is a regular singularity with the characteristic exponents g and $1-g$. In terms of a discrete symmetry transformation $x \to ix$, the eigenfunction with degree v is mapped to a virtual state wavefunction

$$\varphi_{0,\mathrm{v}}(x) \stackrel{\text{def}}{=} e^{x^2/2} x^g L_{\mathrm{v}}^{(g-1/2)}(-x^2), \quad \tilde{\mathcal{E}}_{0,\mathrm{v}} = -4\mathrm{v} - 2g + 3, \quad \mathrm{v} = 0, 1, \ldots, \tag{3.2}$$

in which $L_n^{(\alpha)}(x)$ is the Laguerre polynomial of degree n in x. The nodelessness of $\varphi_{0,\mathrm{v}}(x)$ is obvious, since all the zeros of the Laguerre polynomial $L_n^{(\alpha)}(x)$ are positive. The boundary conditions (A) and (B) are satisfied. The Wronskian* of these virtual state wave functions

$$\mathrm{W}[\varphi_{0,\mathrm{v}_1}, \ldots, \varphi_{0,\mathrm{v}_M}](x) \tag{3.3}$$

can also be used for compositions, as demonstrated for the multi-indexed orthogonal polynomials [3]. The situation is the same for all the other examples below. The radial oscillator itself provides an infinite number of different compositions.

*$\mathrm{W}[f_1, \cdots, f_n](x) \stackrel{\text{def}}{=} \det\left(\partial_x^{j-1} f_k(x)\right)_{1 \le j,k \le n}$.

3.2 Pöschl-Teller

The Pöschl-Teller potential is

$$V_0(x) = \frac{g(g-1)}{\sin^2 x} + \frac{h(h-1)}{\cos^2 x}, \quad 0 < x < \frac{\pi}{2}, \quad g, h > 1/2. \tag{3.4}$$

The lower boundary $x = 0$ is a regular singularity with the characteristic exponents g and $1-g$. The upper boundary $x = \frac{\pi}{2}$ is also a regular singularity with the characteristic exponents h and $1-h$. In terms of a discrete symmetry transformation $h \to 1-h$, the eigenfunction with lower degree v is mapped to a virtual state wavefunction

$$\varphi_{0,\mathrm{v}}(x) \stackrel{\text{def}}{=} (\sin x)^g (\cos x)^{1-h} P_{\mathrm{v}}^{(g-1/2,1/2-h)}(\cos 2x), \tag{3.5}$$

$$\tilde{\mathcal{E}}_{0,\mathrm{v}} \stackrel{\text{def}}{=} (g-h+1+2\mathrm{v})^2, \qquad \mathrm{v} = 0, 1, \ldots, [h-1/2]'. \tag{3.6}$$

Here $P_n^{(\alpha,\beta)}(x)$ is the Jacobi polynomial of degree n in x and $[a]'$ denotes the greatest integer less than a. It is easy to see that the boundary conditions (A) and (B) are satisfied. The expansion of the Jacobi polynomial can be used to demonstrate the nodelessness. (See (3.2) of [16].)

3.3 Hyperbolic Pöschl-Teller

The hyperbolic Pöschl-Teller potential

$$V_0(x) = \frac{g(g-1)}{\sinh^2 x} - \frac{h(h+1)}{\cosh^2 x}, \quad 0 < x < \infty, \quad h > g > 1/2, \tag{3.7}$$

has $[(h-g)/2]' + 1$ discrete eigenstates. The lower boundary $x = 0$ is a regular singularity with the characteristic exponents g and $1-g$. By the discrete symmetry transformation $h \to -(h+1)$, the eigenfunction with degree v is mapped to a virtual state wave function

$$\varphi_{0,\mathrm{v}}(x) \stackrel{\text{def}}{=} (\sinh x)^g (\cosh x)^{h+1} P_{\mathrm{v}}^{(g-1/2,h+1/2)}(\cosh 2x), \tag{3.8}$$

$$\tilde{\mathcal{E}}_{0,\mathrm{v}} \stackrel{\text{def}}{=} -(h-g)^2 - (2\mathrm{v}+2g+1)(2\mathrm{v}+2h+1), \quad \mathrm{v} = 0, 1, \ldots, [(h-g)/2]'. \tag{3.9}$$

It is easy to see that the boundary conditions (A) and (B) are satisfied. The expansion of the Jacobi polynomial can be used to demonstrate the nodelessness.

The *overshoot eigenfunctions* [4] also provides the proper nodeless solutions. They have exactly the same form as the eigenfunctions but their degrees are much higher than that of the highest eigenstate so that their energies are lower than the groundstate energy:

$$\varphi_{0,\mathrm{v}}(x) \stackrel{\text{def}}{=} (\sinh x)^g (\cosh x)^{-h} P_{\mathrm{v}}^{(g-1/2,-h-1/2)}(\cosh 2x), \tag{3.10}$$

$$\tilde{\mathcal{E}}_{0,\mathrm{v}} \stackrel{\text{def}}{=} -(h-g-2\mathrm{v})^2, \quad \mathrm{v} > h-g. \tag{3.11}$$

It is easy to see that the boundary conditions (A) and (B) are satisfied. The expansion of the Jacobi polynomial can be used to demonstrate the nodelessness.

3.4 Rosen-Morse

This potential is

$$V_0(x) \stackrel{\text{def}}{=} -\frac{h(h+1)}{\cosh^2 x} + 2\mu \tanh x, \quad -\infty < x < \infty, \quad h > \sqrt{\mu} > 0. \tag{3.12}$$

The system has finitely many discrete eigenstates $[h-\sqrt{\mu}\,]'+1$. The overshoot eigenfunctions $h < \mathrm{v} < h + \frac{\mu}{h}$ provides proper nodeless solutions:

$$\varphi_{0,\mathrm{v}}(x) \stackrel{\text{def}}{=} e^{-\frac{\mu}{h-\mathrm{v}}x} (\cosh x)^{-h+\mathrm{v}} P_{\mathrm{v}}^{(\alpha_\mathrm{v},\beta_\mathrm{v})}(\tanh x), \tag{3.13}$$

$$\alpha_\mathrm{v} \stackrel{\text{def}}{=} h - \mathrm{v} + \frac{\mu}{h-\mathrm{v}}, \quad \beta_\mathrm{v} \stackrel{\text{def}}{=} h - \mathrm{v} - \frac{\mu}{h-\mathrm{v}}, \tag{3.14}$$

$$\tilde{\mathcal{E}}_{0,\mathrm{v}} \stackrel{\text{def}}{=} -(h-\mathrm{v})^2 - \frac{\mu^2}{(h-\mathrm{v})^2}, \quad h < \mathrm{v} < h + \frac{\mu}{h}. \tag{3.15}$$

It is easy to see that the boundary conditions (A) and (B) are satisfied. The expansion of the Jacobi polynomial can be used to demonstrate the nodelessness.

3.5 Eckart

This potential problem is also called Kepler problem in hyperbolic space:

$$V_0(x) \stackrel{\text{def}}{=} \frac{g(g-1)}{\sinh^2 x} - 2\mu \coth x, \quad 0 < x < \infty, \quad \sqrt{\mu} > g > \frac{1}{2}. \tag{3.16}$$

It has a finite number of discrete eigenstates, $[\sqrt{\mu} - g]' + 1$. The lower boundary $x = 0$ is a regular singularity with the characteristic exponents g and $1-g$. The overshoot eigenfunctions $\mathrm{v} > \frac{\mu}{g} - g$ provide proper nodeless solutions:

$$\varphi_{0,\mathrm{v}}(x) \stackrel{\text{def}}{=} e^{-\frac{\mu}{g+\mathrm{v}}x} (\sinh x)^{g+\mathrm{v}} P_{\mathrm{v}}^{(\alpha_\mathrm{v},\beta_\mathrm{v})}(\coth x), \tag{3.17}$$

$$\alpha_\mathrm{v} \stackrel{\text{def}}{=} -g - \mathrm{v} + \frac{\mu}{g+\mathrm{v}}, \quad \beta_\mathrm{v} \stackrel{\text{def}}{=} -g - \mathrm{v} - \frac{\mu}{g+\mathrm{v}}, \tag{3.18}$$

$$\tilde{\mathcal{E}}_{0,\mathrm{v}} \stackrel{\text{def}}{=} -(g+\mathrm{v})^2 - \frac{\mu^2}{(g+\mathrm{v})^2}, \quad \mathrm{v} > \frac{\mu}{g} - g. \tag{3.19}$$

It is easy to see that the boundary conditions (A) and (B) are satisfied. The expansion of the Jacobi polynomial can be used to demonstrate the nodelessness.

4 Summary and comments

General prescriptions for composing two potentials V_0 and V_1 in one dimensional quantum mechanics are presented by extending the original work in [1]. The virtual state wavefunctions [3] play an important role, as they have played in the rational deformations of solvable potentials [3]–[12]. The form of the mapping function $\psi_0(x)$ (2.7) is reminiscent of the Abraham-Moses transformations [17], [18], [19], or the so-called binary Darboux transformations [20]. The virtual state wavefunctions also play important roles in Abraham-Moses transformations [19].

Acknowledgements

R. S. is supported in part by Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), No.22540186.

Appendix

References

[1] K. Chadan and R. Kobayashi, "New classes of potentials for which the radial Schrödinger equation can be solved at zero energy, II" J. Phys. **A39** (2006) 3381-3396 and 13691-13699.

[2] K. Chadan and P. C. Sabatier, *Inverse problems in quantum scattering theory*, second edition, Springer Verlag, New York (1988).

[3] S. Odake and R. Sasaki, "Exactly solvable quantum mechanics and infinite families of multi-indexed orthogonal polynomials," Phys. Lett. **B702** (2011) 164-170, `arXiv:1105.0508[math-ph]`.

[4] S. Odake and R. Sasaki, "Extensions of solvable potentials with finitely many discrete eigenstates," J. Phys. **A46** (2013) 235205 (15pp), `arXiv:1301.3980[math-ph]`.

[5] D. Gómez-Ullate, N. Kamran and R. Milson, "Two-step Darboux transformations and exceptional Laguerre polynomials," J. Math. Anal. Appl. **387** (2012) 410-418, `arXiv:1103.5724[math-ph]`.

[6] D. Gómez-Ullate, N. Kamran and R. Milson, "An extension of Bochner's problem: exceptional invariant subspaces," J. Approx Theory **162** (2010) 987-1006, arXiv:0805.3376[math-ph]; "An extended class of orthogonal polynomials defined by a Sturm-Liouville problem," J. Math. Anal. Appl. **359** (2009) 352-367, arXiv:0807.3939[math-ph].

[7] C. Quesne, "Exceptional orthogonal polynomials, exactly solvable potentials and supersymmetry," J. Phys. **A41** (2008) 392001 (6pp), arXiv:0807.4087[quant-ph].

[8] B. Bagchi, C. Quesne and R. Roychoudhury, "Isospectrality of conventional and new extended potentials, second-order supersymmetry and role of PT symmetry," Pramana J. Phys. **73** (2009) 337-347, arXiv:0812.1488[quant-ph].

[9] S. Odake and R. Sasaki, "Infinitely many shape invariant potentials and new orthogonal polynomials," Phys. Lett. **B679** (2009) 414-417, arXiv:0906.0142[math-ph].

[10] C. Quesne, "Solvable rational potentials and exceptional orthogonal polynomials in supersymmetric quantum mechanics," SIGMA **5** (2009) 084 (24pp), arXiv:0906.2331[math-ph].

[11] S. Odake and R. Sasaki, "Another set of infinitely many exceptional (X_ℓ) Laguerre polynomials," Phys. Lett. **B684** (2010) 173-176, arXiv:0911.3442[math-ph].

[12] C.-L. Ho, S. Odake and R. Sasaki, "Properties of the exceptional (X_ℓ) Laguerre and Jacobi polynomials," SIGMA **7** (2011) 107 (24pp), arXiv:0912.5447[math-ph].

[13] L. Infeld and T. E. Hull, "The factorization method," Rev. Mod. Phys. **23** (1951) 21-68.

[14] F. Cooper, A. Khare and U. Sukhatme, "Supersymmetry and quantum mechanics," Phys. Rep. **251** (1995) 267-385.

[15] S. Bochner, "Über Sturm-Liouvillesche Polynomsysteme," Math. Zeit. **29** (1929) 730-736.

[16] S. Odake and R. Sasaki, "A new family of shape invariantly deformed Darboux-Pöschl-Teller potentials with continuous ℓ," J. Phys. A **44** (2011) 195203 (14pp), arXiv:1007.3800[math-ph].

[17] P. B. Abraham and H. E. Moses, "Changes in potentials due to changes in the point spectrum: Anharmonic oscillators with exact solutions," Phys. Rev. **A22** (1980) 1333-1340.

[18] M. Luban and D. L. Pursey, "New Schrödinger equations for old: Inequivalence of the Darboux and Abraham-Moses constructions," Phys. Rev. **D33** (1986) 431-436; D. L. Pursey, "New families of isospectral Hamiltonians," Phys. Rev. **D33** (1986) 1048-1055.

[19] S. Odake and R. Sasaki, "Non-polynomial extensions of solvable potentials à la Abraham -Moses," J. Math. Phys. **54** (2013) 102106 (19pp), `arXiv:1307.0931[math-ph]`.

[20] V. B. Matveev and M. A. Salle, *Darboux transformations and solitons*, Springer-Verlag, Berlin Heidelberg (1991).